



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\text{Mass of sphere} = \pi \int_0^a (a^4 - x^4) dx = \frac{4}{5} \pi a^5.$$

$\therefore \frac{4}{5} \pi a^5 = 3\pi(a^4 - \frac{1}{5})$ ,  $4a^5 = 15a^4 - 3$ , which evidently has a root slightly less than 3.75.

In the solution given by Dr. Zerr in the November MONTHLY if the parts be added so as to give the mass of the sphere the result is not homogeneous in  $a$  and is therefore evidently wrong. In getting  $M$  the upper limit for  $\theta$  should be  $\cos^{-1}(1/r)$  and not  $\cos^{-1}(1/a)$ . For  $M_1$  we must subtract  $2M$  from the mass of the sphere.

75. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

A particle  $P$ , is held in a bent tube by two forces directed towards two fixed points,  $H$  and  $S$ . Show that the equation of the tube is  $PS \cdot PH = k^2$ , if the forces are  $\mu/PS$  and  $\mu/PH$ .

I. Solution by GEORGE R. DEANE, C. E., B. S., Professor of Mathematics, Missouri School of Mines, Rolla, Mo.

Put  $PS = r_1$ ,  $PH = r_2$ . By the principle of virtual work, we have,

$$\frac{\mu}{r_1} \delta r_1 + \frac{\mu}{r_2} \delta r_2 = 0.$$

Let  $f(r_1, r_2) = 0$  be the equation of the curve. Then

$$\frac{\partial f}{\partial r_1} \delta r_1 + \frac{\partial f}{\partial r_2} \delta r_2 = 0.$$

Eliminating  $\delta r_1$  and  $\delta r_2$ ,

$$\frac{\frac{\partial f}{\partial r_1}}{\frac{\partial f}{\partial r_2}} = \frac{\frac{\mu}{r_1}}{\frac{\mu}{r_2}}.$$

$$\text{Whence, } -\frac{dr_2}{dr_1} = \frac{r_2}{r_1}, \quad r_1 dr_2 + r_2 dr_1 = 0, \quad r_1 r_2 = k^2.$$

The general theorem of which this is a particular case, is given in Minchin's *Statics*, Vol. I., page 88.

II. Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio State University, Athens, Ohio.

Let  $s$  = any arc of the tube,  $r, r'$  = the distances of the particle from the centers of force at any time  $t$ ,  $m$  = the same absolute intensities of the forces, and  $\beta$  = the velocity of projection.

If  $S, S'$  be the radial forces attracting the particle, we will have

$$\frac{d^2 s}{dt^2} = -S \frac{dr}{ds} - S' \frac{dr'}{ds} \dots \dots (1).$$

But  $S=m/r$ ,  $S'=m/r'$ , and (1) becomes

$$\frac{d^2 s}{dt^2} = -\frac{m}{r} \frac{dr}{ds} - \frac{m}{r'} \frac{dr'}{ds} \dots \dots (2).$$

Multiply by  $2(ds/dt)$  and integrate ; then

$$\frac{ds^2}{dt^2} = -m \log r^2 - m \log r'^2 + C \dots \dots (3).$$

When  $r=a$ ,  $r'=a$ ,  $\frac{ds}{dt}=\beta$ ;  $\therefore C=\beta^2-m^2 \log \frac{1}{a^4}$ , and (3) is

$$\beta^2 = m^2 \log \frac{1}{r^2 r'^2} + \beta^2 - m^2 \log \frac{1}{a^4} \dots (4),$$

or,  $rr'=a^2 \dots \dots (5)$ , a lemniscate.

[Other solutions of this problem will appear in the next issue.]

## DIOPHANTINE ANALYSIS.

73. Proposed by M. A. GRUBER, A. M., War Department, Washington, D. C.

Find integral values for  $x$  and  $y$  in  $\left( \begin{matrix} 2x^2 - y^2 = \square \\ 2y^2 - x^2 = \square \end{matrix} \right)$ .

I. Solution by the PROPOSER.

$$2x^2 - y^2 = \square = a^2 \dots \dots (1), \quad 2y^2 - x^2 = \square = (2).$$

From (1),  $y^2 = 2x^2 - a^2$ . Substituting this in (2), we have  $3x^2 - 2a^2 = b^2$ . Whence  $x = \frac{1}{3}\sqrt{[3(2a^2 + b^2)]}$ , and  $y = \frac{1}{3}\sqrt{[3(a^2 + 2b^2)]}$ .

As far as I know, the only method of rationalizing both radicals is to put  $a=b$ . Then  $x=y=a=b$ .

Accordingly, no *different* integral values can be found for  $x$  and  $y$ .

This problem is the key to Problem 62, "To find four squares in arithmetical progression."

The roots of the four squares would then be, respectively,

$$a, \quad \frac{1}{3}\sqrt{[3(2a^2 + b^2)]}, \quad \frac{1}{3}\sqrt{[3(a^2 + 2b^2)]}, \quad b.$$

The common difference of the squares is  $\frac{1}{9}(b^2 - a^2)$ .

According to the above solution, the roots of the four squares could not all be rational integers ; *one* of them, at least, must be a *surd*. It is evident, however, that an infinite number of sets of four squares can be found in which *two* of the roots are rational integers.